

UNITARY REPRESENTATIONS OF THE FUNDAMENTAL GROUP OF ORBIFOLDS

INDRANIL BISWAS AND AMIT HOGADI

ABSTRACT. We generalize to smooth orbifolds the correspondence between the polystable vector bundles and unitary representations for a smooth projective variety.

1. INTRODUCTION

Let $Y/\mathbb{C}$ be a connected smooth projective curve and $\mathcal{E} \rightarrow Y$ a polystable vector bundle of degree zero. A celebrated theorem of Narasimhan and Seshadri, [5], says that $\mathcal{E}$ is necessarily given by a unitary representation of the fundamental group of Y . Let $X/\mathbb{C}$ be a smooth projective variety of dimension n . Fix an ample line bundle $\mathcal{L}$ on X in order to define the degree of a torsionfree coherent sheaf on X . The following generalization of the Narasimhan–Seshadri theorem holds.

1.1. Theorem ([4], [1]). *Let $\mathcal{E}$ be a vector bundle on X such that $c_1(\mathcal{E}) = 0$ and $c_2(\mathcal{E}) \cdot c_1(\mathcal{L})^{n-2} = 0$. Then $\mathcal{E}$ is polystable if and only if it is given by a unitary representation of the (topological) fundamental group of X .*

Our aim here is to generalize the above theorem to projective orbifolds. By an **orbifold** over a field k we always mean a smooth separated Deligne Mumford stack which is of finite type over k .

1.2. Theorem. *Let $X/\mathbb{C}$ be an irreducible projective orbifold of dimension n , and let $\mathcal{L}$ be an ample line bundle on X . Let $\mathcal{E} \rightarrow X$ be a vector bundle (with respect to $\mathcal{L}$) on X such that $c_1(\mathcal{E}) = 0$ and $c_2(\mathcal{E}) \cdot c_1(\mathcal{L})^{n-2} = 0$. Then $\mathcal{E}$ is polystable if and only if it is obtained by a unitary representation of $\pi_1^{\text{top}}(X, x)$ for a (equivalently, any) closed point $x \in X$.*

Here $\pi_1^{\text{top}}(X, x)$ denotes the topological fundamental group of underlying complex analytical stack. The above theorem follows quite easily from Theorem 1.1 in the special case when X is a global quotient of a smooth variety by a finite group. The main work needed to prove Theorem 1.2 is to deduce the general case from this special case. As a consequence of Theorem 1.2, we obtain the validity of Theorem 1.1 certain types of singular varieties.

2000 *Mathematics Subject Classification.* 14F05, 14J60.

Key words and phrases. Orbifolds, polystable vector bundle, unitary representation.

1.3. Corollary. *Let $X/\mathbb{C}$ be a projective variety with at worst quotient singularities. Assume X is the coarse moduli space of a projective orbifold $\tilde{X}/\mathbb{C}$ (e.g. X has isolated singular points). Then Theorem 1.1 holds for X .*

Proof. Let $\mathcal{L}$ be an ample line bundle on X and $\mathcal{E}$ be a polystable vector bundle satisfying $c_1(\mathcal{E}) = 0$ and $c_2(\mathcal{E}) \cdot c_1(\mathcal{L})^{n-2} = 0$ where n is the dimension of X . Let $\pi : \tilde{X} \rightarrow X$ be the given morphism. By 1.2, $\pi^*(E)$ is given by a unitary representation of $\pi_1^{top}(\tilde{X}, x)$ (for some geometric point x of $\tilde{X}$). However, since the vector bundle arising from this representation is a pull back from X , it is clear that for every point y of $\tilde{X}$, the isotropy group G_y at y lies in the kernel of this representation. Thus by using ([6], 8.1) this descends to give a unitary representation of $\pi_1^{top}(X, x)$, which defines the vector bundle $\mathcal{E}$. $\square$

2. RIGIDIFICATION

Let X be a Deligne Mumford stack over $\mathbb{C}$, and let x be a geometric point of X . Throughout we use the notation $\pi_1^{top}(X, x)$ to denote the topological fundamental group of the underlying complex analytic stack, and use $\pi_1(X, x)$ to denote the algebraic fundamental group.

Recall that the notion of rigidification of a Deligne Mumford stack is a process to get rid of given stabilizers in a “minimal” manner.

2.1. Definition. Let $f : Y \rightarrow X$ be a 1-morphism of locally noetherian Deligne Mumford stacks. f is called a **rigidification** if

- (1) f is separated and of finite type, and
- (2) for any atlas $U \rightarrow X$, where U is a scheme, the projection $Y \times_X U \rightarrow U$ is a coarse moduli space.

2.2. Remark. Since the morphism to the coarse moduli space of a separated Deligne Mumford Stack is always proper, it follows that a rigidification morphism is always a proper morphism.

Let $f : Y \rightarrow X$ be a morphism of connected noetherian schemes. Let y be a geometric point of Y , and let $x = f(y)$. For any geometric point y' of Y , let $G_{y'}$ denote the isotropy group at y' . The kernel of the homomorphism

$$G_{y'} \rightarrow G_{f(y')}$$

will be denoted by $K_{y'}$. There is a group homomorphism

$$G_{y'} \rightarrow \pi_1(Y, y)$$

well defined up to conjugation by elements of $\pi_1(Y, y)$. Let $N(f)$ denote the closed normal subgroup of $\pi_1(Y, y)$ generated by images of the homomorphisms

$$(1) \quad K_{y'} \rightarrow \pi_1(Y, y),$$

where y' runs over all geometric points of Y .

When Y is a stack of finite type over $\mathbb{C}$, there are also homomorphisms

$$G_{y'} \longrightarrow \pi_1^{\text{top}}(Y, y)$$

and one defines $K_{y'}^{\text{top}}$ to be the kernel of the above map and $N^{\text{top}}(f)$ to be the normal subgroup of $\pi_1^{\text{top}}(Y, y)$ generated by images of the above homomorphisms for all $y' \in Y(\mathbb{C})$.

2.3. Theorem. *Let $f : Y \longrightarrow X$ be a proper morphism of connected noetherian Deligne Mumford stacks. Let y be a geometric point of Y , and let $x = f(y)$. Assume that there exists a dense open substack $U \subset X$ such that the codimension of $X \setminus U$ is at least two, and the induced morphism*

$$f|_U : f^{-1}(U) \longrightarrow U$$

is a rigidification. Then the following two statements hold:

(i) *The sequence of groups*

$$1 \longrightarrow N(f) \longrightarrow \pi_1(Y, y) \longrightarrow \pi_1(X, x) \longrightarrow 1$$

is exact.

(ii) *If X and Y are schemes of finite type over $\mathbb{C}$ and $y \in Y(\mathbb{C})$, then the following similar sequence of the topological fundamental groups*

$$1 \longrightarrow N^{\text{top}}(f) \longrightarrow \pi_1^{\text{top}}(Y, y) \longrightarrow \pi_1^{\text{top}}(X, x) \longrightarrow 1$$

is exact.

Proof. We will prove (i); the proof of (ii) is similar. Let us first consider the case when f is a rigidification. Let $Et(Y)$ denote the category of finite étale coverings of Y . Let $\tilde{\mathcal{C}}$ denote the full subcategory of objects $Z \xrightarrow{h} Y$ in $Et(Y)$ such that for any geometric point z of Z ,

$$K_{h(z)} \subset \text{Image}(\pi_1(Z, z) \longrightarrow \pi_1(Y, h(z))).$$

We claim that the category $Et(Y)$ is equivalent to the category $Et(X)$ of finite étale coverings of X . If $X' \longrightarrow X$ is a finite étale covering, then it is easy to see that $X' \times_X Y \longrightarrow Y$ is in $Et(Y)$. Conversely, let $Z \longrightarrow Y$ be a finite étale covering in $\tilde{\mathcal{C}}$. Then we need to show that there is a natural finite étale cover $X' \longrightarrow X$ such that $Z = X' \times_X Y$. This can be proved étale locally on X and hence we may assume that X is in fact the coarse moduli space of Y in which case this is simply the result of [6].

Now consider the general case where $f : Y \longrightarrow X$ is a rigidification in codimension one. Let $U \subset X$ be the open subset of X whose complement has codimension at least two and such that $f^{-1}(U) \longrightarrow U$ is a rigidification. In this case, the proof follows from the above special case and the following well known equivalence of categories.

$$Et(Y) \cong Et(f^{-1}(U)) \cong Et(U) \cong Et(X).$$

This completes the proof of the theorem. $\square$

2.4. Theorem. *Let k be any field of characteristic zero. Let X/k be any quasiprojective orbifold. Then there exists a 1-morphism $\phi : Y \longrightarrow X$ such that*

- (1) *$\tilde{Y}$ is an orbifold which is a finite global quotient, and*
- (2) *There exists a dense open subset $V \subset X$ such that $\phi^{-1}(V) \longrightarrow V$ is a rigidification morphism and the complement of V has codimension at least two.*

Proof. Let $D \subset X$ be a (reduced) divisor such that q is an isomorphism over $X \setminus D$. Since X is normal, there exists an open subset $V \subset X$ such that

- (1) $D \cap V$ is smooth.
- (2) $X \setminus V$ has codimension at least two.

Let m be a positive integer such that for every point p of X , the order of the isotropy group at p divides m . Let $\mathcal{L}$ be a sufficiently ample line bundle on X and let D' be a section of $\mathcal{L}^m \otimes \mathcal{O}_X(-D)$ such that $(D + D') \cap V$ is smooth. Note that since X is not smooth, D may not be a Cartier divisor hence the reflexive sheaf $\mathcal{O}_X(-D)$ may not be a line bundle. We replace D by $D + D'$ without loss of generality and assume that there exists a line bundle $\mathcal{L}$ on X such that $\mathcal{L}^m \cong \mathcal{O}_X(D)$.

Let T denote the cyclic cover of X defined by the section D of $\mathcal{L}^m$. There is a natural $G = \mathbb{Z}/m$ action on T given by the action of $\mathbb{G}_m$ on $\mathcal{L}$; by the choice of the integer m we have a rational map $T \dashrightarrow X$ defined over all codimension one points of $\tilde{X}$. Now we 'resolve' the singularities of the rational map $T \dashrightarrow X$ G -equivariantly, to get a proper birational G -map $T' \longrightarrow T$ such that T' is smooth and the induced map $T' \longrightarrow X$ is actually a morphism. Moreover, by construction, this map is G -invariant. Thus we have an induced morphism from $\phi : Y \longrightarrow X$ where $Y = [T'/G]$. It is now straight-forward to check that the map ϕ satisfies the required conditions of the theorem. $\square$

3. POLYSTABLE VECTOR BUNDLES ON ORBIFOLDS

In this section we prove Theorem 1.2. Throughout this section we work over the field $k = \mathbb{C}$.

3.1. Definition. Let Z be a projective Deligne Mumford Stack over k . A line bundle $\mathcal{L}$ on Z is called ample, if some power of $\mathcal{L}$ descends as an ample line bundle on the coarse moduli space of Z .

3.2. Lemma. *Let X/k be a projective orbifold, and let $\mathcal{L}$ be an ample line bundle on X . Let $f : Y \longrightarrow X$ be a finite morphism, where Y/k is a normal projective variety. Let $\mathcal{E}$ be a vector bundle on X . Then*

- (i) *$\mathcal{E}$ is semistable with respect to $\mathcal{L}$ if and only if $f^*\mathcal{E}$ is semistable w.r.t. $f^*(\mathcal{L})$.*
- (ii) *$\mathcal{E}$ is polystable with respect to $\mathcal{L}$ if and only if $f^*\mathcal{E}$ is polystable with respect to $f^*(\mathcal{L})$.*

Proof. The proof is similar to the case when X is a variety instead of an orbifold, and we refer the reader to [2, 3.2] for details of the argument. $\square$

Let X and $\mathcal{L}$ be as above. Let n be the dimension of X .

3.3. Proposition. *Let $f : Y \longrightarrow X$ be any morphism of projective orbifolds over k . Let $\mathcal{E}$ be a polystable vector bundle on X such that $c_1(\mathcal{E}) = 0$, and $c_2(\mathcal{E}) \cdot c_1(\mathcal{L})^{n-2} = 0$. Then $f^*\mathcal{E}$ is polystable vector bundle on Y .*

Proof. By [3] we can find a diagram

$$\begin{array}{ccc} Y' & \xrightarrow{h} & Y \\ \downarrow f' & & \downarrow f \\ X' & \xrightarrow{g} & X \end{array}$$

where Y' and X' are smooth projective varieties and the horizontal morphisms are finite and dominant. By Lemma 3.2, the pullback $g^*\mathcal{E}$ is polystable, and also, $c_1(g^*\mathcal{E}) = 0$, and $c_2(g^*\mathcal{E}) \cdot c_1(g^*\mathcal{L})^{n-2} = 0$; note that $g^*\mathcal{L}$ is ample because g is finite. Thus, $g^*\mathcal{E}$ is given by a unitary representation of the fundamental group of X' [1], [4]. Therefore, $f'^*g^*\mathcal{E}$ is also given by a unitary representation of the fundamental group of Y' , and hence it is polystable. Thus by Lemma 3.2, $f^*\mathcal{E}$ is polystable. $\square$

3.4. Remark. We first observe that proving the Theorem 1.2 is equivalent to showing that if $\mathcal{E} \longrightarrow X$ is polystable, and $\pi_X : U_X \longrightarrow X$ is the universal covering space of X , then

- (1) $\pi_X^*(\mathcal{E})$ is trivial, and
- (2) $\pi_X^*(\mathcal{E})$ admits a $\pi_1^{\text{top}}(X, x)$ invariant flat unitary metric.

3.5. Lemma. *Theorem 1.2 holds if X is a finite global quotient.*

Proof. If $\mathcal{E} \longrightarrow X$ is given by a unitary representation of $\pi_1^{\text{top}}(X, x)$, then using Theorem 1.1, Lemma 1.1 and Proposition 3.3 it follows that $\mathcal{E}$ is polystable, and $c_1(\mathcal{E}) = 0 = c_2(\mathcal{E}) \cdot c_1(\mathcal{L})^{n-2} = 0$.

Assume that $\mathcal{E} \longrightarrow X$ is a polystable vector bundle with $c_1(\mathcal{E}) = 0 = c_2(\mathcal{E}) \cdot c_1(\mathcal{L})^{n-2} = 0$. By hypothesis, there exists a smooth projective variety Y and a finite group G acting on Y such that $X = [Y/G]$. Let $f : Y \longrightarrow X$ denote the natural map. Then $f^*(\mathcal{E})$ is a polystable bundle on Y with $c_1(f^*\mathcal{E}) = 0$ and $c_2(f^*\mathcal{E}) \cdot c_1(f^*\mathcal{L})^{n-2} = 0$. Let $\pi_Y : U_Y \longrightarrow Y$ denote the universal covering space of Y . Then $\pi_Y^*f^*(\mathcal{E})$ is a trivial vector bundle on U_Y with $\pi_1^{\text{top}}(Y, y)$ invariant unitary flat metric, say $\langle -, - \rangle$. Note that U_Y is also the covering space for X . Moreover if $x = f(y)$, then $\pi_1^{\text{top}}(Y, y)$ is naturally a finite index subgroup of $\pi_1^{\text{top}}(X, x)$. Let $\{g_i\}$ denote its finite coset representatives. It is then clear that the “averaged” metric,

$$(v_1, v_2) := \sum_i g_i^* \langle v_1, v_2 \rangle$$

is a $\pi_1^{top}(X, x)$ invariant unitary flat metric on U_Y . $\square$

We now reduce the general case of Theorem 1.2 to the above special case using Theorem 2.4.

Proof of Theorem 1.2. As in Lemma 3.5, if $\mathcal{E}$ is given by a unitary representation of $\pi_1^{top}(X, x)$, then $\mathcal{E}$ is polystable, and $c_1(\mathcal{E}) = 0$ and $c_2(\mathcal{E}) \cdot c_1(\mathcal{L})^{n-2} = 0$.

Assume that $\mathcal{E}$ is polystable, and $c_1(\mathcal{E}) = 0$ and $c_2(\mathcal{E}) \cdot c_1(\mathcal{L})^{n-2} = 0$. Let $\phi : Y \rightarrow X$ be a morphism as guaranteed by Theorem 2.4. Fix a point y_0 of Y . Then, since Y is a finite global quotient, by Lemma 3.5, $\phi^*(\mathcal{E})$ is defined by a unitary representation of $\pi_1^{top}(Y, y_0)$.

However, for any point $y \in Y$, the kernel of $G_y \rightarrow G_{\phi(y)}$ acts trivially on the fiber of $\phi^*(\mathcal{E})$ at y . Thus this kernel is also contained in the kernel of the unitary representation. Then by Theorem 2.3, the unitary representation of $\pi_1^{top}(Y, y_0)$ which defines $\phi^*(\mathcal{E})$ actually factors through $\pi_1^{top}(X, \phi(y_0))$. This completes the proof. $\square$

REFERENCES

- [1] S.K. Donaldson, Infinite determinants, stable bundles and curvature. *Duke Math. Jour.* **54** (1987), 231–247.
- [2] D. Huybrechts, M. Lehn, *The geometry of moduli spaces of sheaves*. Aspects of Mathematics, **E31**. Friedr. Vieweg & Sohn, Braunschweig, 1997.
- [3] A. Kresch, A. Vistoli, On coverings of Deligne-Mumford stacks and surjectivity of the Brauer map. *Bull. London Math. Soc.* **36** (2004), 188–192.
- [4] V. Mehta, A. Ramanathan, Restriction of stable sheaves and representations of the fundamental group. *Invent. Math.* **77** (1984), 163–172.
- [5] M.S. Narasimhan, C.S. Seshadri, Stable and unitary vector bundles on a compact Riemann surface. *Ann. of Math.* **82** (1965), 540–567.
- [6] B. Noohi, Fundamental groups of algebraic stacks. *Jour. Inst. Math. Jussieu*, **3** (2004), 69–103.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400005, INDIA

E-mail address: indranil@math.tifr.res.in

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, BOMBAY 400005, INDIA

E-mail address: amit@math.tifr.res.in